

इंटरनेट

मानक

Disclosure to Promote the Right To Information

Whereas the Parliament of India has set out to provide a practical regime of right to information for citizens to secure access to information under the control of public authorities, in order to promote transparency and accountability in the working of every public authority, and whereas the attached publication of the Bureau of Indian Standards is of particular interest to the public, particularly disadvantaged communities and those engaged in the pursuit of education and knowledge, the attached public safety standard is made available to promote the timely dissemination of this information in an accurate manner to the public.

“जानने का अधिकार, जीने का अधिकार”

Mazdoor Kisan Shakti Sangathan

“The Right to Information, The Right to Live”

“पुराने को छोड़ नये के तरफ”

Jawaharlal Nehru

“Step Out From the Old to the New”


IS 11816-3 (1986): Storage of cereals and pulses, Part 3:
Control of attack by vertebrate and invertebrate animals
[FAD 16: Foodgrains, Starches and Ready to Eat Foods]



“ज्ञान से एक नये भारत का निर्माण”

Satyanarayan Gangaram Pitroda

“Invent a New India Using Knowledge”



“ज्ञान एक ऐसा खजाना है जो कभी चुराया नहीं जा सकता है”

Bhartrhari—Nitiśatakam

“Knowledge is such a treasure which cannot be stolen”

BLANK PAGE



IS : 11816 (Part 3) - 1986

(Reaffirmed 2005)

Indian Standard

STORAGE OF CEREALS AND PULSES

**PART 3 CONTROL OF ATTACK BY VERTEBRATE AND
INVERTEBRATE ANIMALS**

UDC 633'1+635'65 : 664'8'03

© Copyright 1987

**BUREAU OF INDIAN STANDARDS
MANAK BHAVAN, 9 BAHADUR SHAH ZAFAR MARG
NEW DELHI 110002**

Indian Standard

STORAGE OF CEREALS AND PULSES

PART 3 CONTROL OF ATTACK BY VERTEBRATE AND INVERTEBRATE ANIMALS

The Storage Structures and Storage Management Sectional Committee, AFDC 28

Chairman

DR K. KRISHNAMURTHY

Representing

Department of Food (Ministry of Food and
Civil Supplies), New Delhi

Members

- | | |
|---|---|
| DR G. K. GIRISH (<i>Alternate to</i>
Dr K. Krishnamurthy) | |
| LT-COL K. N. ACHARYA | Army Headquarters (QMG's Branch) (Ministry
of Defence), New Delhi |
| LT-COL S. K. BHATTACHARYA (<i>Alternate</i>) | |
| AGRICULTURAL MARKETING
ADVISER TO THE GOVERNMENT OF
INDIA | Directorate of Marketing and Inspection
(Ministry of Agricultural and Rural
Development), Faridabad |
| SHRI P. S. AMBERMANI (<i>Alternate</i>) | |
| DR A. ALAM | Indian Council of Agricultural Research, New
Delhi |
| PROF A. P. BHATNAGAR | Punjab Agricultural University, Ludhiana |
| DR MALVINDER SINGH (<i>Alternate</i>) | |
| DR C. D. DWIVEDI | Defence Materials and Stores Research and
Development Establishment, Kanpur |
| SHRI V. K. MATHUR (<i>Alternate</i>) | |
| SHRI K. PANDITHURAI | Food Corporation of India, New Delhi |
| SHRI K. S. BHASIN (<i>Alternate</i>) | |
| SHRI M. V. PAVATE | Planning Commission, Government of India,
New Delhi |
| SHRI S. K. SINGH (<i>Alternate</i>) | |
| SHRI G. PERUMAL | Central Public Works Department, New Delhi |
| PLANT PROTECTION ADVISER TO
THE GOVERNMENT OF INDIA | Directorate of Plant Protection, Quarantine
and Storage, Faridabad |
| SHRI T. R. SHARMA (<i>Alternate</i>) | |
| SHRI K. S. N. RAJU | National Co-operative Development Corpora-
tion, New Delhi |
| SHRI R. L. GAUR (<i>Alternate</i>) | |
| DR SONE LAL | Indian Grain Storage Institute, Hapur |
| SHRI B. R. BIREWAR (<i>Alternate</i>) | |

(Continued on page 2)

© Copyright 1987

BUREAU OF INDIAN STANDARDS

This publication is protected under the *Indian Copyright Act* (XIV of 1957) and reproduction in whole or in part by any means except with written permission of the publisher shall be deemed to be an infringement of copyright under the said Act.

(Continued from page 1)

Members

Representing

SHRI P. R. VISHWAMBHARAM	Central Warehousing Corporation, New Delhi
SHRI AJAY KHERA ((Alternate I)	
SHRI A. P. MATHUR (Alternate II)	
DR T. D. YADAV	Indian Agricultural Research Institute (ICAR), New Delhi
SHRIMATI S. SINGH (Alternate)	
SHRI T. PURNANANDAM, Director (Agri & Food)	Director General, BIS (Ex-officio Member)

Secretary

SHRIMATI SHASHI SAREEN
Deputy Director (Agri & Food), BIS

Storage Management Subcommittee, AFDC 28 : 9

Convener

DR K. KRISHNAMURTHY	Department of Food (Ministry of Food and Civil Supplies), New Delhi
DR G. K. GIRISH (Alternate to Dr K. Krishnamurthy)	

Members

LT-COL K. N. ACHARYA	Army Headquarters (QMG's Branch), (Ministry of Defence), New Delhi
LT-COL S. K. BHATTACHARYA (Alternate)	
SHRI V. S. AGARWAL	Indian Produce Association, Calcutta
DR S. S. BAINS	Punjab Agricultural University, Ludhiana
DR B. S. CHAHAL (Alternate)	
SHRI R. N. DEWAN	Punjab State Co-operative Supply and Marketing Federation Limited, Chandigarh
CIVIL ENGINEER (Alternate)	
DIRECTOR (FOOD & SUPPLIES)	Food and Supplies Department, Government of Punjab, Chandigarh
SHRI S. S. BEDI (Alternate)	
DR C. D. DWIVEDI	Defence Materials and Stores Research and Development Establishment, Kanpur
SHRI V. K. MATHUR (Alternate)	
SHRI R. P. JAIN	Roller Flour Millers Federation of India, New Delhi
SHRI RAGHUNANDAN BHATIA (Alternate)	
SHRI S. K. MAJUMDAR	Central Food Technological Research Institute (CSIR), Mysore
MANAGER (STORAGE)	Food Corporation of India, New Delhi
SHRI MANGALDAS NATHUBHAI	Grain, Rice and Oilseeds Merchants' Association, Bombay
PLANT PROTECTION ADVISER TO THE GOVERNMENT OF INDIA	Directorate of Plant Protection, Quarantine and Storage, Faridabad
SHRI D. SRINATH (Alternate)	
DR M. S. PRADHAN	National Cooperative Development Corporation, New Delhi
SHRI J. P. MITTAL (Alternate)	
SHRI ADUSUMILLI SUBHA RAO	Andhra Pradesh Millers' Association, Vijaywada
SHRI A. C. SHROFF	Excel Industries Ltd, Bombay
SHRI P. V. KANGO (Alternate)	
DR SONE LAL	Indian Grain Storage Institute, Hapur
SHRI B. R. BIREWAR (Alternate)	

Indian Standard

STORAGE OF CEREALS AND PULSES

PART 3 CONTROL OF ATTACK BY VERTEBRATE AND INVERTEBRATE ANIMALS

0. FOREWORD

0.1 This Indian Standard (Part 3) was adopted by the Indian Standards Institution on 30 May 1986, after the draft finalized by the Storage Structures and Storage Management Sectional Committee had been approved by the Agricultural and Food Products Division Council.

0.2 This standard is being published in the following three parts:

Part 1 General consideration in keeping cereals

Part 2 Essential requirements

Part 3 Control of attack by vertebrate and invertebrate animals.

0.3 In the preparation of this standard (Part 3), guidance has been taken from ISO 6322/3-1981 'Storage of cereals and pulses: Part 3, Control of attack by vertebrate and invertebrate animals' issued by the International Organization for Standardization.

1. SCOPE

1.1 This standard (Part 3) gives guidance on means of controlling attack by vertebrate and invertebrate animals on cereals and pulses in storage.

1.1.1 Other aspects of storage of cereals and pulses are dealt with in Parts 1 and 2 of this standard.

2. VERTEBRATE PESTS

2.1 General

2.1.1 The ability of birds, rats, mice and other rodents to feed on grain is not limited by the quality or condition of the grain but only by its accessibility.

2.1.2 The effects of attack by birds, rats, mice and other rodents are contamination, spillage, loss in mass and lowering of quality. Contamination may, for example, prevent wheat from being used for flour milling because of excessive count of rodent hairs in the finished flour.

IS : 11816 (Part 3) - 1986

2.1.3 Attention to hygiene and proofing of stores against access are the principal preventive measures, although anticoagulant baits (such as warfarin) may also be used for routine control of rats and mice. The use of an acute poison may be necessary if resistance to anticoagulants appears in rat or mouse populations.

2.1.4 Birds should be excluded from grain stores by stringent bird-proofing by fixing wire gauge steel frames on doors, windows, ventilators and other exits. Outside the stores, birds may be scared away or repelled by suitable means.

2.2 Control

2.2.1 Rats and mice may be controlled by trapping, or by the use of a bait incorporating a poison which acts within a few minutes of ingestion (acute poison) or by one which acts over a period (chronic poison) or by fumigation with gas. Fumigation is generally used as a part of an integrated control system, as it normally gives no residual protection.

2.2.2 Fumigation against rodents is normally performed at lower dosage rates and for shorter periods of exposure than those required for insect control. It follows that an insect fumigation also controls rodents. Methods involving fumigation or the use of acute poisons should be carried out only by properly trained persons, as specified by governmental regulations. Some poisons and fumigants, which may be used, are listed in Appendix A.

3. INVERTEBRATE PESTS

3.1 General

3.1.1 Attack by insects and mites is influenced by the accessibility of cereals and pulses and also by the type and condition of the grain.

3.1.2 Attack may not only lead to loss in mass and to contamination, but may alter the condition and quality of the grain.

3.1.3 Infestation of grain both by true storage pests and by field crop pests, may result in the contamination of milled products with insect fragments.

3.2 Onset of Infestation

3.2.1 Pulses

3.2.1.1 Pulses are liable to be attacked during growth by various moths and pulse beetles (Lepidoptera and Coleoptera) [for example, *Bruchus pisorum* (L.) (pea beetle)] which lay eggs on the developing pods. The larvae penetrate into the seeds and continue to develop in the dry seeds after harvest, adult pulse beetles eventually emerging and

flying to the fields to start the cycle again. Some species [for example, *Callosobruchus maculatus* (F.) (cowpea beetle) and *Acanthoscelides obtectus* (Say.) (dried bean beetle)] are capable of continuing to reproduce on the dry peas and beans. Seeds containing pupae or adults ready to emerge can be recognized by the thin outer layer of the seed coat, or ' window ', covering the end of the exit tunnel. Seeds from which beetles have emerged have clean round holes. Tests for infestation of pulses are described in IS : 6261-1971*.

3.2.1.2 The storage of pulses should therefore be organized on the assumption that they are already infested when taken into store. The risk depends on local circumstances (climate, field control methods, etc), being greatest in the tropics, where conditions favour survival and continuous rapid development of pests.

3.2.2 Cereals

3.2.2.1 Field infestation of cereals by storage insects occurs mainly in the tropics and sub-tropics but also in the warmer parts of temperate regions. It is normally caused by insects flying out from storage buildings to lay eggs on ripening crops. Some improved varieties of hybrid maize with short sheath leaves surrounding the cob are particularly liable to such infestation. Insects causing such infestation are mainly *Sitotroga cerealella* (Oliv.) (Angoumois grain moth) and *Sitophilus zeamais* (Motsch) (greater maize weevil).

3.2.2.2 In most temperate climates, infestation by storage insects usually starts after the harvested crop has been placed in store, but may commence immediately if the combine harvester or sacks harbour insect pests.

3.2.2.3 Storage of cereals in areas where field infestation by storage insects may occur should take this risk into account. Storage of cereals in most temperate areas can be organized on the basis that grain from the field is free from infestation, except for certain species of mites.

3.2.2.4 Some species of mites carried with grain from the field can continue to breed in store (for example, *Tyrophagus longior* Gerv.) but, more commonly, the dominant mites are true storage species [for example, *Acarus siro* L. (flour mite)].

3.2.2.5 Mites are a particularly grave problem wherever grains are stored under high atmospheric relative humidities or their moisture contents exceeds 15 percent (m/m), even when the temperatures are too low for insect development.

*Methods of analysis for detection of insect and rodent contamination in grains and milled products.

3.3 Factors Affecting Insect and Mite Development

3.3.1 A few species of insects [for example, *Sitophilus granarius* (L.) (grain weevil)] can attack completely sound grain, but abrasion of the seed coat, breakage during harvesting and movement and admixtures of cereal dust, facilitate attack by many other species of insects and mites [for example, *Oryzaephilus surinamensis* (L.) (saw toothed grain beetle), *Acarus siro* L. (flour mite)].

3.3.2 The insects and mites found in stored cereals include those which tunnel in the endosperm [for example, *Sitophilus granarius* (L.) (grain weevil), *Sitophilus oryzae* (L.) (rice weevil)], those which consume the germ [for example, *Cryptolestes ferrugineus* (Steph.) (rust red grain beetle), *Plodia interpunctella* (Hubn.) (Indian meal moth) and *Acarus siro* L. (flour mite)], those which attack broken pieces and dust [for example, *Tribolium castaneum* (Herbst.) (rust red flour beetle), *Glycyphagus destructor* (Schr.) (forage mite)], those which live on moulds [for example, fungus beetles (Cryptophagidae and Mycetophagidae)], those which are parasites or predators on other insects and mites [for example, Braconidae and Cheyletidae) and those which are scavengers on dead bodies of other insects (for example, Dermestidae). Many species are capable of performing more than one role [for example, *Tenebroides mauritanicus* (L.) (cadelle beetle), a predator and an occasional feeder on germ].

3.3.3 The principal factors which affect the development and activity of insect and mite pests are:

- a) temperature,
- b) moisture content of the grain,
- c) relative humidity of the immediate atmosphere, and
- d) nature of the grain.

3.3.3.1 Most insect pests of stored grain cannot complete the full development from egg to adult, mating and further egg production at temperatures below 10°C or greater than 35°C. The minimum temperature required for the multiplication of most of the serious insect pests is about 15°C and the lower end of the optimum range is at least 20°C. For most mites, the limits and optima are generally 5°C lower.

3.3.3.2 The moisture content of grain affects insects and mites directly since they absorb water with their food and indirectly through controlling the relative humidity of the inter-granular air. For each species, there is a minimum relative humidity below which complete development cannot take place or which is lethal. Grain of less than 9 percent (m/m) moisture content (equilibrium relative humidity of 30 percent) is generally secure from insect or mite attack, although

Trogoderma granarium Everts (khapra beetle) can breed at a moisture content of 2 percent.

3.3.3.3 For each species, there is a combination of temperature and humidity at which populations increase most rapidly. This means that different species tend to occur in those parts of India where the climate is most suitable for them.

3.3.3.4 A summary of these conditions for the most common species of grain pests is given in Appendix B.

3.4 Heating of Grain Caused by Insects and Mites

3.4.1 Large bulks or bagged stacks of cereals and pulses tend to stabilize the conditions of temperature and humidity within them. It is these conditions, in particular, which directly affect the insects, and not the diurnal fluctuations in the free space of the store. Local variations in the temperature and moisture contents of bulks are exploited by insects and mites which tend, by random movement, to find their way to favourable breeding places. If conditions here are above the minima shown in Appendix B, the insects will breed. As they develop they produce heat which may not disperse as rapidly as it is produced. Temperature rises and insect development accelerates. Eventually the temperature rises to an unfavourable level, usually not above 42°C, and those insects which can move outwards do so; those which cannot (for example, larvae within grains) eventually die. The living insects finally concentrate at the surface.

3.4.2 Temperature gradients are established in the bulk, and moisture moves from the hot interior to the cooler periphery. If the temperature at the surface is below the dew point of the moist air from the interior, condensation will take place and the grain may eventually sprout. Moisture content and equilibrium relative humidity may be raised locally to a point at which the growth of moulds is initiated. Thus, in grain nominally safe for definite storage, 'dry grain heating', initiated by insects, may change to 'damp grain heating' caused by micro-organisms. The temperature in 'damp grain heating' seldom rises above 62°C and usually remains at about 52°C.

3.5 Prevention and Control of Insect and Mite Infestation

3.5.1 Prevention

3.5.1.1 Attack may be prevented by denying access to insects and mites, by maintaining an environment unfavourable for breeding, or by applying a protective treatment to the grain for seed purposes only. The principal techniques for prevention are hygiene, that is the removal of all unwanted grain and dust in which insects and mites can breed,

treatment of empty locations, and control of the environment, by such means as:

- a) keeping grain below the temperature or the humidity necessary for increase in pest numbers (for example, drying, aerated storage, and refrigerated storage);
- b) hermetic or inert-atmosphere storage; and
- c) enclosing pest-free grain in insect-resistant containers.

3.5.2 Control

3.5.2.0 Control may include killing or removing insects and mites:

- a) in empty storage buildings,
- b) in empty transport vehicles,
- c) in containers (including sacks) before they are filled with grain, and
- d) in the grain itself.

3.5.2.1 *Mechanical and physical methods* — Grain may be screened to remove free living insects and mites; it can be treated with inert dusts; it can be heated (but care is needed because there is only a small differential between the temperature needed to kill insects and that at which grain may be damaged) : treatment with gamma radiation, and other high frequency electromagnetic waves of accelerated electrons are being used in many countries; it can be placed in airtight storage, in which insects are killed when they reduce the oxygen content in the atmosphere to below 2 percent.

3.5.2.2 *Biological methods* — There are very few circumstances in which biological control (that is, the use of predators, parasites or pathogens) is likely to be effective in the control of cereal pests, because the levels of stabilized populations are normally too high to be tolerated. The bacterium *Bacillus thuringiensis* may be used commercially for the control of moths in stored wheat, maize, sorghum grain, rough rice and soya beans.

3.5.2.3 *Chemical methods* — Chemicals are used in two main forms: contact insecticides (some pesticides in general use for treatment of empty stores and of seed grain only are listed in Appendix A) and fumigants.

- a) *Contact insecticides* — Residual contact insecticides and acaricides applied as dusts, oil-in-water emulsions or water dispersible sprays, aerosols or smokes — render the surfaces of empty stores, transport vehicles, ships, cargo containers and conveying equipment toxic to insects and mites. When the insects and mites cross these surfaces they pick up sufficient toxic material to kill them before they reach the commodity to be protected.

This technique is effective against crawling insects and mites. The period for which the insecticide persists is dependent on the nature of the surface, temperature, moisture content of the air and extent of exposure to light. Flying insects may be killed by the use of aerosols or smokes whilst in flight, or when they alight on a treated surface; flying insects and those on the surface of commodities may also be controlled by volatile insecticides intermediate in properties between contact insecticides and fumigants.

- b) *Fumigants* — Control of pest populations deep in the structure of buildings or vehicles, in empty sacks and in the grain itself, whether in bags or bulk, can be achieved only by fumigants, which act as gases, even though they may be applied as solids or liquids, and which normally have no residual effect on insects or mites after the fumigant has dispersed. Fumigant gases are generally toxic to man and should only be used by properly trained persons, who know the dangers and the necessary safeguards.
- c) *Pesticide residues* — In India the sale and use of insecticides is regulated by the Government whose requirement keeps in mind the possibilities of contamination of any grain to which they may be applied. The recommendations of the FAO/WHO Codex Alimentarius Commission regarding limits for residues in food are taken into account in such regulations. Only those insecticides which are specifically approved in this way should be used in grain stores and even then they should be used strictly in accordance with the manufacturer's recommendations and any Government requirements in regard to safety in use. The recommendation of the Inter-Governmental Maritime Consultative Organization (IMCO) regarding the use of pesticides in ships should also be taken into account. Such use should also take into account the requirements of purchasers (for example, some maltsters require for barley a certificate of freedom from malathion) within a country and in other countries.

A P P E N D I X A
(*Clauses 2.2.2 and 3.5.2.3*)

**SOME PESTICIDES USED FOR TREATMENT OF GRAIN,
EMPTY GRAIN STORES AND VEHICLES**

A-1. INSECTICIDES

A-1.1 Contact Insecticides (for Seed Purposes only)

- a) *Pyrethroids*: Pyrethrins or synthetic pyrethrins with a synergist.
- b) *Organochlorine Compounds*: gamma-HCH (gamma-BHC, lindane). Organochlorine compounds are now prohibited almost all over the world.
- c) *Organophosphorous Compounds*: Only those approved by Central Insecticides Board, Ministry of Agriculture and Rural Development.

A-1.2 Fumigants

- a) Methyl bromide.
- b) Hydrogen phosphide.
- c) Liquid bulk fumigants, pure or as mixtures (for example ethylene dichloride, ethylene dibromide, carbon tetrachloride).
- d) Chloropicrin (only as an admixture of methyl bromide).

A-1.2.1 Only properly trained persons shall be employed to use fumigants.

A-2. RODENTICIDES

A-2.1 Acute poisons and burrow fumigants, for example, zinc phosphide, barium carbonate and aluminium phosphide.

Only properly trained person shall be employed to use acute poisons.

A-2.2 Chronic poisons, such as hydroxycoumarins (for example, warfarin, coumafuryl, coumachlor).

APPENDIX B

(Clauses 3.3.3.4 and 3.4.1)

LIMITING AND OPTIMUM CONDITIONS FOR INCREASE OF POPULATIONS OF CERTAIN INSECT AND MITE PESTS OF STORED PULSES AND CEREALS

SPECIES	COMMON NAME	MINIMUM		OPTIMUM		MAXIMUM RATE OF MULTIPLICATION PER MONTH (x)	SUSCEPTIBILITY OF PRODUCTS TO ATTACK					
		Temperature, °C	Relative Humidity, Percent	Temperature, °C	Relative Humidity, Percent, Minimum		C	C*	C**	c	P	p
<i>Acanthoscelides obtectus</i> (Say.)	dried bean beetle	17	30	27 to 31	65	25					*	
<i>Acarus siro</i> L.	flour mite	7	65	21 to 27	80	2 500		*		*	*	*
<i>Callosobruchus chinensis</i> (L.)	cowpea beetle	19	30	28 to 32	60	30					*	
<i>Callosobruchus maculatus</i> (F.)	cowpea beetle	22	30	30 to 35	50	50					*	
<i>Corcyra cephalonica</i> (Staint.)	rice moth	18	30	28 to 32	30	10		*		*		
<i>Cryptolestes ferrugineus</i> (Steph.)	rust red grain beetle	23	10	32 to 35	65	60		*		*		
<i>Cryptolestes pusillus</i> (Schön.)	flat grain beetle	22	60	28 to 33	70	10		*		*		
<i>Ephestia cautella</i> (Wlk.)	tropical warehouse moth	17	25	28 to 32	60	50		*		*		
<i>Ephestia elutella</i> (Hübn.)	warehouse moth	10	30	25	70	15		*		*		*
<i>Ephestia kuehniella</i> Zeller	Mediterranean flour moth	10	1	24 to 27	65	50		*		*		
<i>Lasioderma serricorne</i> (F.)	cigarette beetle	22	30	32 to 35	55	20				*	*	
<i>Laheticus oryzae</i> Waterh.	long headed flour beetle	26	30	33 to 37	—	10			*	*		

(Continued)

SPECIES	COMMON NAME	MINIMUM		OPTIMUM		MAXIMUM RATE OF MULTIPLICATION PER MONTH (x)	SUSCEPTIBILITY OF PRODUCTS TO ATTACK					
		Temperature, °C	Relative Humidity, Percent	Temperature, °C	Relative Humidity, Percent, Minimum		C	C*	C**	c	P	p
<i>Oryzaephilus surinamensis</i> (L.)	saw toothed grain beetle	21	10	31 to 34	65	50		*		*		
<i>Plodia interpunctella</i> (Hübner.)	Indian meal moth	18	40	28 to 32	—	30		*		*		
<i>Ptinus tectus</i> Boield.	Australian spider beetle	10	50	23 to 25	70	4		*		*		*
<i>Rhizopertha dominica</i> (F.)	lesser grain borer	23	30	32 to 35	50	20	*			*		
<i>Sitophilus granarius</i> (L.)	grain weevil	15	50	26 to 30	70	15	*					
<i>Sitophilus oryzae</i> (L.)	rice weevil	17	60	27 to 31	70	25	*					
<i>Sitotroga cerealella</i> (Oliv.)	Angoumois grain moth	16	30	26 to 30	—	50	*					
<i>Tribolium castaneum</i> (Herbst.)	rust red flour beetle	22	1	32 to 35	65	70			*	*		*
<i>Tribolium confusum</i> Duv.	confused flour beetle	21	1	30 to 33	50	60			*	*		*
<i>Trogoderma granarium</i> Everts.	khapra beetle	24	1	33 to 37	45	12.5	*			*	*	*
<i>Zabrotes subfasciatus</i> (Boh.)		22	30	29 to 33	50	20					*	

C — Whole undamaged cereals, C* — Germ of cereals, C** — Cereal grain already damaged mechanically or by other insects, c — Cereal products, P — Whole pulses, p — Pulse products.